

CA2AFRICA

Working document

Implementation of Conservation Agriculture in the highlands of Vakinankaratra, Madagascar; constraints and opportunities.

A non-adoption case study

Maaïke Hartog (Wageningen),

Eric Penot (CIRAD UMR Innovation) &

Yan de Graff (Wageningen)

2011

1 INTRODUCTION

Madagascar, *la Grande Île*, is a place on earth with some most peculiar natural features. The current degraded status of its soils is unfortunately less unique. The vulnerable environment suffers from the increasing pressure of the growing population. Cropping on slopes causes erosion. Soil fertility is low, and soil nutrient inputs are needed. The concept of Conservation Agriculture (CA) has been introduced as the way to fight worldwide soil degradation and change soil mining agriculture into a more sustainable system. At this time, the concept is used on 6-7% of the world's cropland (FAO, 2009). The adoption rate among African smallholders has been very limited. This is the starting point for the EU-project CA2AFRICA. It aims at examining the conditions that determine success or failure of CA (CA2AFRICA, 2009). The project uses three scales to analyze CA: field, farm/village and regional. This research is focused on the scale of farm and village. This research focuses on the highlands in the region of Vakinankaratra. NGO's and scientists have undertaken activities to introduce CA, but so far the practices have not been implemented on a large scale in the highlands. The research has been carried out from June to September 2010.

2 RESEARCH CONTEXT

REGIONAL INFORMATION

RESEARCH LOCATION

The research takes place in Madagascar, an island-nation situated in the Indian Ocean. Vakinankaratra is a region of Madagascar (see figure 2-1). The choice for the study area was made on the basis of the administration of the BVPI-project, that defines its zones according to the irrigation zones (*perimètres irrigués*). For this research two zones were chosen: Fitakimerina and landratsay (both part of *Lot 1* of the project). This was done because of their proximity to the town of Antsirabe, and because of the low adoption rate of CA in these zones. There is a third zone that falls into this category: Ikabona, but it was not chosen because of the limited time. Note that these zones (*perimètres irrigués*) do not correspond with the administrative units that carry the same name.




Figure 2-1: Vakinankaratra (red) and study areas (landratsay and Fitakimerina), Source: CIRAD

After the region, the next administrative level is that of districts, which are divided into *communes*. The research is conducted in the *communes* Vaninkarena (district Antsirabe II) and Mandritsara (district Ikabona). The level that can be identified as 'village' is called *fokontany*. Because of the complicated structure and the difference between BVPI and Malagasy administration, this report refers to locations and sublocations rather than *fokontany* or county. The following sublocations have been visited:

Location	Sublocation	Number of interviews
Fitakimerina (<i>commune</i> Vaninkarena)	Anjanamanjaka (BVPI: Ambohimanga&Ambohitraivo) Tsaratanana	26 4
landratsay (<i>commune</i> Mandritsara)	Ambohimarina (BVPI: Est Anosy) Ampamelomana	20 10

NATURAL FEATURES

The highlands of Vakinankaratra have a warm rainy season from October/November to March/April. The winter season is dry and cold, from April/May to September. Figure 2-2 gives a general picture of the precipitation in this region. Table 2.1 presents natural features per location.




Figure 2-2 Monthly rainfall for 2003-2009, station Andranomalenatra-Kobama (Source: BVPI SE/HP, 2009b)

Table 2.1 Natural features of study locations

Location	Sublocation	Altitude (meter above sea level)	Soil types	Annual precipitation
Fitakimerina	Anjanamanjaka		Hills: Acid ferralsol (pH 5.0), texture: sandy loam, poor in OM, very poor in P,Ca,Mg and K. Valleys: Deposit of lakes and rivers.	1200 mm, mainly in January and February
	Tsaratanana	1500	Tsaratanana: Occurrence of hydromorph soil at flat <i>tanety</i> .	
landratsay	Ambohimarina		Hills : Rich volcanic and poor ferralsol. Valleys : Less rich.	1450 mm in 110 days (85% in November to March)
	Ampamelomana	1600*		

*) Estimation from GoogleEarth
Source: BVPI SE/HP, 2008a-c

SOCIO-ECONOMIC FEATURES

In 2008, 3 million tonnes of rice were produced in Madagascar. From the 1.9 million tonnes in 1970, this figure has steadily grown, as has the population of *la grande île*. The food supply per capita has decreased since then and was about 160 kg /year in 2008 (FAOSTAT, 2010). Agricultural policy has mostly been directed towards the intensification of rice cultivation. Since the eighties of last century, a persistent economic crisis withheld farmers from buying the inputs needed for this intensive cultivation. This led to more diversification, by growing crops during the counter season and making more use of *tanety* fields (Seguy, 2003). The city of Antsirabe plays an important role in this region. The highlands around the city supply food, especially dairy products, to its 159 000 inhabitants. The dairy sector is sensitive to political struggle, since governors also have their share in economical activities. Antsirabe is home to the brewery of STAR. This company buys barley from the farmers through an organisation called Malto. Farmers receive the right seeds and inputs, money that is extracted from the price they get for the harvested barley. Especially in landratsay, this is an important source of income. At the studied locations, agriculture is the main economic activity. Three different seasons can be distinguished:

- Main season: From October (transplanting rice from nursery to fields) until March/April/May (harvest of rice)
- Intermediate season at *tanety* (hillsides): From December (after beginning of the rains)
- Counter season at valleys: From March/April/May (harvest of rice) until October

During the intermediate season, most households have consumed all of their rice. This is the period in which most people buy rice, even if they have been selling rice at harvest time (at a lower price).

An important feature of the area is the difference between the *tanety* and the valleys. Figure 3 shows a flat valley, with barley growing in the paddy fields during the counter season. The *tanety* surrounding it are mostly terraced.



Figure 2-3 Typical landscape for the region, photographed in landratsay.

It is not always possible to irrigate and drain the paddy fields effectively. Some fields are qualified as RMME : *rizières à mauvaise maîtrise de l'eau*, paddy fields that are not effectively drained or

irrigated. In this kind of fields, the transplanting of rice from the nursery happens relatively late. In landratsay, some fields are inundated during the rainy season (BVPI SE/HP, 2008a-c).

In table 2.2, some information on population and farming systems can be found per study location.

Table 2.2 Socio-economic features of study locations

Location	Sublocation	Inhabitants	Presence of cows	Principal crops	
				Tanety	Paddy's counter season
Fitakimerina	Anjanamanjaka	2000 (Ambohimanga)	25% of inhabitants possesses Zebu cattle.	In order of importance : Maize, beans, pluvial rice, sweet potato, cassava, bambara groundnut.	In order of importance : Potato, barley, wheat, peas, garlic.
	Tsaratana	700			
landratsay	Ambohimarina	725*	65% of inhabitants possesses Zebu, average of 3/household. Dairy cattle sporadic.	Main season: Maize, beans, soja, potato, rice.	Wheat, barley, taro, beans, maize for cows, tomato, ray grass, legumes, peas, potato.
	Ampamelomana	Not known			

*) Information from president of *fokontany*, 31-07-2010

Source: BVPI SE/HP, 2008a-c

The tenure status of most land is not legalized. At the location of Fitakimerina, 2/3 of the land is bought and 1/3 inherited. 50% of the farmers are hiring the land. (BVPI SE/HP, 2008a) In landratsay, most of the people are owners, but without a formal title (BVPI SE/HP, 2008b). Apart from agriculture, the inhabitants have some opportunities to gain an off-farm income. In Ambohimanga (Fitakimerina), sand is mined from the river. This activity is most important in the winter season, just like the fabrication of bricks. Other sectors include work with the electricity plant of Jirama (the electricity company) and construction of buildings (BVPI SE/HP, 2008a). Several people also run a small shop or restaurant, or have a small revenue from handicraft from wheat stalks (BVPI SE/HP, 2008b).

Table 2.3 Farm data by (sublocation, from BVPI-database of Vakinankaratra, end of August 2010

Location	Sublocation	Selected BVPI parameters	Number of persons (p)	Area exploited/p (ha)	Average total figures			
					Area paddy/p (ha)	No. draft oxen	No. Zebus	No. dairy cows
Fitakimerina	Anjanamanjaka	<i>Villages</i> Ambohimanga & Ambohitraivo	51	3.05	0.88	0.84	2.18	0.61
	Tsaratana	<i>Village</i> Tsaratana	34	1.22	0.22	0.44	0.64	0.12
landratsay	Ambohimarina	<i>Terroir</i> Est Anosy	102	1.03	0.27	2.24	1.73	1.64
	Ampamelomana	<i>Terroir</i> Ampamelomana	100	1.44	0.57	2.33	4.23	1.99

Source: BVPI SE/HP, 2010a

CONSERVATION AGRICULTURE

The soils of the world are degrading and lose their fertility, partly as a result of conventional tillage practices. Negative effects of tillage are for example oxidation of organic matter (OM) and destruction of pores. The uncovered state of the soil after ploughing increases erosion risks (Hobbs, 2007). Several authors (Hobbs, 2007; Fowler & Rockström, 2001) have indicated the need to make agriculture more sustainable. Sustainability is generally defined as a way of acting that does not "... compromise the ability of future offspring to produce their food needs by damaging the natural resources used to feed the population today" (Hobbs *et al.*, 2008). The alternative agricultural practices that are being developed, were by the Food and Agriculture Organisation of the United Nations (FAO) considered as a package, and labelled as 'Conservation Agriculture'. These practices are: Continuous minimum mechanical soil disturbance, Permanent organic soil cover & Diversification of crop species grown in sequence or associations.(FAO, 2010)

It aims at making better use of the agricultural resources in order to minimize external inputs. The continuous soil cover protects the surface structure, so that the soil can keep more water. In a brochure by the French Development Agency (AFD, 2007), a difference is made between CA and direct seeding mulch-based cropping systems. The latter term refers to cropping systems that exclude tillage and have permanent plant cover on the soil - be it with a living crop or dead mulch. In French, these techniques are called: *Système de Culture sur Couverture Végétale*. The concept of direct seeding mulch-based cropping systems was launched by CIRAD in 1999. It does not include crop rotation explicitly, but it still can be seen as a special form of CA that does not differ much from the FAO definition. In this proposal, the term 'CA practices' is consistently used to refer to this particular package.

APPLICABILITY OF CA PRINCIPLES

The promotion of Conservation Agriculture cannot take place without a critical reflection on the principles of these practices. If we take the plough as a metaphor, the paradigm shift that took place has changed this object from a useful tool to a dangerous soil-destroyer. Gowing and Palmer (2008) call it a paradigm shift because CA requires a way of thinking that puts the health of the soil ecosystem first. From this way of thinking, tillage becomes disturbance of the soil and has very damaging effects. The new paradigm includes a shift from soil mining agriculture to sustainable agriculture. Advocates of CA underline the importance of producing more food from less resources (Hobbs *et al.*, 2008). The confidence of the FAO in promoting CA becomes clear in citations like: "The cases where CA did not perform as expected can usually be related to mistakes or shortcuts in the management of the system, but not to any inherent failures in the system" (FAO, 2009). In a paper by Giller *et al.* (2009), it is suggested that CA can be inappropriate in many cases. It places a heavy burden on the livestock feeding, since crop residues are normally used for fodder. Another issue is the shift of labour from ploughing (men's work) to hand weeding (mostly done by women). The same article points to the fact that CA is brought as a 'inseparable' package. It is not clear what the effect of mulching is, when practiced apart from the minimum-tillage. Rabary *et al.* (2008) admit that their study was also unable to isolate this effect. However, they add that mulching and conventional tillage is a combination that is not commonly practiced. According to Wall (2007), small-scale farmers are generally prepared to invest in a profitable CA-system, but lack access to (especially initially) necessary inputs like herbicides and fertilizers. CA also implies changes in agricultural management and the planning of a crop. This makes it a knowledge intensive system. Smallholders are

disadvantaged because their linkages are often within the community and less oriented to outside information networks. To know whether the principles of CA are applicable for this particular situation, the highlands of Madagascar, the pros and cons need to be profoundly analyzed. The following chapters will elaborate on this.

ADOPTION OF CONSERVATION AGRICULTURE

The process of adoption has been studied since the first extension services encountered their first disappointing results. Adoption here is defined as the phase where a farmer makes effort and invests in the implementation of conservation agriculture. De Graaff *et al.* (2008) distinguish between three phases: acceptance, actual adoption and continued use. This last phase is the term that I use for implementing CA without intensive supervision/sponsoring by a project. It indicates that farmers are intrinsically motivated to maintain and replicate the CA measures. The final phase can only be achieved when farmers have experience with the measures, it requires time.

Many different factors can be thought of when studying the behaviour of farmers as they decide about adoption or non-adoption. But it is important to note that it is not a black-or-white decision. The intensity of implementation is important to distinguish between trials and actual adoption. Farmers also do not adopt a blueprint that is determined for them, they often adapt the measures to their own situation (de Graaff *et al.*, 2008). A research on farmer innovation in East Africa concluded that an increase in income is the most important motivation to innovate. In this context, the author poses that investments in conservation measures are stronger motivated by gains in production (read: income, standard of living) than by environmental concerns (Critchley, 1999). The research of Clay *et al.* (1998), in the highlands of Rwanda, found that the following features were positively related to investments in land conservation: land that is owned, has a medium slope, is less fragmented and is cultivated for a shorter time, and among smaller farmers and those with little land in fallow, woodlot, and pasture. An off-farm income also stimulates these investments, because it provides money, especially when households have little access to credit. The same can be said about cash cropping. Lastly, this research mentions public investments in extension and roads as a way to promote sustainable intensification.

Research in Bolivia and Peru observed that investment in soil- and water conservation (SWC) measures was higher with increasing farm size. Also, market-oriented farmers invested more. In Ethiopia, farmer's age, farm size and steep slopes positively affected the initial use of SWC measures. The involvement in or number of livestock had a significantly negative influence on it. But farm size was negatively correlated to continued use, along with participation in off-farm work and family size. In an overall comparison of these independent research projects, only age of head of household and programme participation had a more or less general positive influence on initial adoption. Profitability of SWC measures is mentioned as an important factor for their continued use (de Graaff *et al.*, 2008).

Apart from these resources, a lot more research has been done on the factors that influence farmers' decisions. Knowler and Bradshaw (2007) wrote a synthesis of recent research related to adoption of CA. The analyses that they compared, covered a range of economic, social, physical and institutional factors. The authors conclude that "there are few if any universal variables that regularly explain the adoption of conservation agriculture". The observation that the results of these analyses are so inconsistent, points to the importance of local management.

3 Problematic

It seems that the implementation of CA practices reaches a deadlock in the highland of Vakinankaratra (BVPI SE/HP, 2010b). The objective of the current research is to know why the farmers in this region are not using the CA practices that were suggested to them.

What are the past and present reasons why farmers are hesitant to adopt (certain) CA practices in the highlands of Vakinankaratra?

Sub-questions

1. What were the most important roles of the different development organisations (including research) in the highlands of Vakinankaratra in the past, concerning the adoption of CA practices?
2. What are the past and present reasons* why farmers are hesitant to adopt (certain) CA practices in the highlands of Vakinankaratra? (with special attention to adoption of the particular system that has potential for the area)

*Two groups of reasons are distinguished: Those that can be influenced by farmers and those that cannot (circumstances like climate). These two things strongly interact with each other. The first sub-question is meant to provide background information and will not be treated very extensively. The choice of the location has been made by E. Penot, according to the BVPI project demand and the other research activities concerning CA evaluation in other areas. URP SCRID (*Unité de Recherche en Partenariat 'Systèmes de Culture et Rizicultures Durables'*) is still searching for CA systems adapted to local conditions. This research hopes to contribute to that quest. This study focuses on the process that led to the current situation, both from the side of the extension organisations and the side of farmers. It will not go into details about technical aspects of the measures. Some attention will also be given to the gender aspect of transferring CA practices.

4 METHODOLOGY

Part of the research objective is to understand how CA practices have been introduced from the side of development organisations. I have looked for the opinions of researchers and personnel of the organisations that have experience in the region. The other part is about the farmers. The intention was to start from the point of view of a farmer instead of reasoning from a package of measures that should be implemented. This is why the term 'constraints' is avoided. The survey aimed at interviewing the person who made the decision about CA, be it the man or the woman of the family.

The empirical part of the research consisted of a survey undertaken among 60 farmers. The fieldwork took place during July and August 2010. Preparation and (part of) analysis was done in Antsirabe. During the interviews I stayed in the villages. This paragraph will explain the way the sample was determined.

Randrianarison *et al.* (2008) created a typology with six different types for the *fokontany* Antsapanimahazo:

1. Farmers that have never tried CA;

2. Farmers who abandoned CA after 1 or 2 years;
3. Farmers who abandoned CA after 3 or 4 years;
4. Farmers who abandoned CA after more than 5 years;
5. Farmers practising CA for 1 or 2 years;
6. Farmers practising CA for 3 years or more.

In this typology, farmers of type 5 are called ‘experimental adopters’ and those of type 6 ‘real adopters’. Type 4 was created because these farmers have distinguished reasons for abandoning. Most farmers in the *fokontany* belong to type 1.




Figure 4-1: Percentages of adoption/abandoning of CA in relation to years practiced, in Antsapanimahazo (Source: Randrianarison, 2008)

If farmers have never been into contact with CA practices, they have not yet been in a position to choose. These farmers are not very useful to this research. The CA systems that are introduced in this region, are described in section 5.1 and 5.2. The first level of determining the sample consists of three groups:

1. Those that have never tried CA;
2. Those that have abandoned CA after some time;
3. Those that are still practising CA.

For the sample to be representative, it was subsequently drawn in a stratified random way, according to the strata in table 4.1. The strata of farm size, number of dairy cows and gender of farmer were considered important because they had the highest potential to influence attitude towards CA (according to the theoretical framework). For each group, the research has aimed at the sample in table 4.2. Following from the relatively high percentages of ‘unknown’ in the BVPI-database, the reality was different from these data. The area of the exploitations turned out to be much smaller than expected; in the whole study only one exploitation of more than 4 ha was encountered.

Table 4.1 Farm features, by location, according to BVPI-database of Vakinankaratra, 10 June 2010

	Total	Farm < 4 ha	Size unknown	>1 dairy cow *	Unknown number cows*	Women	Gender unknown
Number Fitakimerina	89	68	14	14	19	32	7
Percentage Fitakimerina	100	76	16	16	21	36	8
Number landratsay	370	307	49	73	281	69	48
Percentage landratsay	100	83	13	20	76	19	13

*) In French: *vache laitière*

Source: BVPI SE/HP, 2010a

Table 4.2 Composition of sample per study location

Farm size	Dairy cows	Gender	Fitakimerina	landratsay
≥ 4 ha	> 0	M/F	1	1
< 4 ha	> 0	M	1	1
< 4 ha	> 0	F	1	1
< 4 ha	0	M	4	5
< 4 ha	0	F	3	2
Total			10	10

All the interviewed persons were selected by the extension agent (technician) at the village. A list from BVPI with persons who practise CA was used to compose the first group. For the two other groups, the agent had a free choice.

FARM TYPOLOGY USED

In 2010, a farm typology (Ahmim-Richard *et al.*, 2010) was made for the regions of Vakinankaratra and Amoron'i Mania (neighbours the former in the south). The results of the current survey will refer to this typology. Appendix A gives the determination keys for categorizing farmers.

Table 4.3 Farm typology for Vakinankaratra region, presented by Ahmim-Richard *et al.*, 2010

		Substantial off-farm income		No substantial off-farm income				
				Self-sufficient in rice		Not self-sufficient in rice		
						Complementary off-farm income	No complementary off-farm income	
Agriculture primary activity	Cattle	1A: cows	1BC: other	4A: cows	4BC: other		6	7A: cows
		No cattle	2		5			
							9: no crop sales	
Agriculture secondary activity		3						

Table 4.3 depicts the structure of this farm typology. It consists of three basic farm types:

- Farmers receiving substantial off-farm income (1-3);
Abbreviated as 'Off'

- Full-time farmers, self-sufficient in rice (4&5);
Abbreviated as 'Prod'
- Resource-poor farmers, not self-sufficient in rice and with little off farm income (6-9).
Abbreviated as 'Poor'

PERSONAL STORIES I

Solofson got disappointed by CA

Solofson Randrianjarivo (52 years old) lives in Anjanamanjaka, Fitakimerina. His wife is teacher at the primary school. Because of her salary, they fall into the category of 'receiving substantial off-farm income' (see typology in 4.3). They have five children, of which two still live with them. They own about 15 are of paddy fields and 12 are at *tanety*. For work at the fields they hire a lot of labour each year.

Solofson tried the CA system one time, but was disappointed with the production of the crops (maize and beans). He knows several groups of people who also have abandoned CA practices.



Figure 4-2 Solofson Randrianjarivo and his wife, with Rarivo (co-researcher)

Razafindrakoto has plans to start with CA

Razafindrakoto (50) is a full-time farmer and self-sufficient in rice. He owns two male Zebu for traction, no cows. He and his wife (43) have four sons. The eldest (22) quit school after primary school and works at the farm; the others are at secondary school. The family lives in Ambohimarina, landratsay.

They own 0.7 hectare of land and use the counter season to grow forage on the paddy fields. Barley is cultivated for Malto. Razafindrakoto has an individual attitude and has not joined a producers' organisation. He is into contact with a technician from SDMAD, who has provided him with seeds of vetch. He has installed it and plans to plant upland rice in it this year. In his opinion, CA only works for people who are very motivated. He would apply it to grow fodder and increase the soil quality by the cover.

Charles has abandoned CA after one year

Charles Rakotondranaivo (59) is an enterprising farmer from Ambohimarina, landratsay. He lives with his wife (50) and 2 sons, of which the eldest has finished secondary school and already has its own parcels and cows. Charles owns 2 hectare of land, 4 Zebus, 2 cows and other cattle. He produces milk and also grows forage. He is active within producer's associations.

Has talked to people from BVPI and explained that it is necessary for him to use the oats as forage for his cows. The system does not work with the little cover that remains; weeds get a lot of space and light to grow. A better opportunity for CA could arise if there comes a cover crop that is not eaten by cows.



Figure 4-3 Entrance to the house of Charles Rakotondranaivo

Volatine practices CA with help from her brother

Volatine Aronjahine Razafindrazaka (23) married five years ago and has two little daughters. This keeps her mostly busy at the house. Her husband (26) works at both their farm and other farms; in winter he also has an income from the production of bricks. The young family lives at Tsaratanana, Fitakimerina. Their farm area is 0.12 hectare and their only livestock consists of some poultry; they belong to the group of resource-poor farmers.

Since 2007, they practice CA on some of their land. Volatine is a sister of the SDMAD-technician in Tsaratanana and he assists them with the techniques. She mentions the positive effect of CA on the soil fertility.

5 Theoretical framework

This chapter aims to answer the question: What are the most important factors that explain the hesitant attitude of farmers towards CA in this region? The first paragraph contains information from the organisations that are active in dissemination of CA, to give an oversight of the current state of affairs. Section 5.2 describes the technical constraints and opportunities while in 5.3 an attempt is made to distillate the farmers' point of view from literature. The chapter finishes with a conclusion, that provides hypotheses to be tested by the survey.

The NGO Tafa was founded on January 1, 1995. It took over the role of 'Blé Kobama', an organisation that had gone bankrupt. This organisation had started working on CA practices (literally: *système de culture avec couverture permanente des sols et technique de semis direct*) since 1991, on an experimental farm at Andranomanelatra. When Tafa took over, a start was made with the dissemination of these new techniques. Since 1995, a partnership was formed with FOFIFA, the national research centre for rural development at Antananarivo University (Tafa-FIFAMOR, 1995). Also, a national network of institutions was set up, called GSDM: Direct seeding group of Madagascar. Since 2004, GSDM coordinates the project 'Support for the dissemination of agro ecological techniques in Madagascar'. This project is funded by AFD and the Malagasy government.

In 2006, the project BVPI SE/HP was started; its full name is: *Développement des Bassins Versants et Périmètres Irrigués dans le Sud Est / Hauts Plateaux*. It is part of the national program of BVPI, a policy that was defined by the government. Bidder is the Minister of Agriculture (*Ministère de l'agriculture, de l'élevage et de la pêche*). The project is mainly funded by AFD (BVPI SE/HP, 2009a-b).

In the past, the BVPI SE/HP project has worked on the infrastructure for irrigation. Operators in the field are SDMAD (*Semis Direct Madagascar*) and BEST. If farmers are part of a producer's association, they get seeds and other inputs which they have to pay back after the harvest. SDMAD is a private company. BVPI first pays the company for the inputs and after the payback of the farmers, SDMAD gives the money again to BVPI. Each region has a technical agent, they are employees of SDMAD. BEST is responsible for the water user associations, that group practically all farmers that use the

paddy fields. In landratsay (Ambohimarina and Ampamelonana) and Anjanamanjaka, the promotion of CA practices started in 2006. Tsaratanana followed in 2007. The scientific part of the project is carried out by CIRAD and FOFIFA. Since 2001, they are combined in one group : URP SCRID (CIRAD, 2010). TAFE has an experimental station in Andranomanelatra (district of Antsirabe II) (Rabary *et al.*, 2008). In Fitakimerina and landratsay, It is not the purpose of TAFE to act at farmers' level. They have a test field at Tsaratanana (Fitakimerina) but that is for forage, not to demonstrate CA practices.

Box 1: Description of cover crops introduced in the region by BVPI

- Brachiaria (*Brachiaria*, 6 different species). Newly introduced, farmers discovered that it could be used as forage, since then exploited for Zebu.
- Crotalaire (*Crotalaire*, different species). Weak production of biomass
- Hairy vetch (*Vicia villosa*). Ued for cattle
- Oats (*Avena sativa*). Used for cattle
- New introduction: Eleusine (*Eleusine coracana*) (not yet known in the area, same can happen as to Brachiaria?), radish, cowpeas

The first diagnostic research in the region was performed in the era of Blé Kobama. It showed that the performance of agricultural activities on *tanety* with acid soils was very poor, with low production levels and labour-intensive practices (Seguy, 2003).

EXPERIENCES FROM BVPI SE/HP

As described above, the project BVPI SE/HP has recently started the introduction of CA practices at the study locations. The CA systems that are currently used by farmers who are part of the project:

Fitakimerina	landratsay
Beans + Oats	
Maize + Beans +Oats (landratsay: + Potato)	
Beans + Brachiaria	Potato +oats
Cassava + Brachiaria	Potato + Wheat
Pois de terre+Brachiaria	Potato +Vetch (low part)
Pluvial/non-irrigated Rice + Crotalaire	Ray-grass + Vetch
Soja + Brachiaria	Barley + Vetch
Soja + Crotalaire	Beans + Vetch (mainly C2/C3)
Brachiaria/Oats pure	Wheat + Vetch

In the zone of Fitakimerina, the dissemination of CA practices has not been successful until now. Since the beginning of the project, the cover crops have been removed from the fields; often not with a direct purpose for fodder but to sell the crop residues or exchange it for fertilizers. This happens because the farmers cannot afford chemical fertilizers and also do not own enough cattle. Farmers also prioritize the rice paddies above the *tanety*. According to BVPI SE/HP reporting, adoption of CA practices cannot be expected in this zone (Raharison & Andrianaivolala, 2009).

In landratsay, the pressure on crop residues is also high. The stalks of the maize are for example used as firewood. But there is a potential for systems that improve the 3-cropping system that is practiced on the *tanety*. In this rotation, oats can be added to provide extra biomass. It will be explained in the next paragraph (Raharison & Andrianaivolala, 2009).

TECHNICAL CONSTRAINTS AND OPPORTUNITIES

The high altitude of this zone limits the choice for cover crops. The cover crops that are mostly used, do not support the cold temperatures. There are not that many leguminosae that grow enough biomass and at the same time are not edible for cattle. If soils are more poor, it takes more time to restructure the soil with a cover crop. In combination with the constraints coming from agricultural practice, this can become a problem (Raharison & Andrianaivolala, 2009).

THE OATS-BASED SYSTEM

The most common system that is practiced on *tanety* in landratsay, consists of two cycles (table 5.1): Maize and beans in the main season, and potatoes, barley or wheat after the harvest of the beans. BVPI SE/HP proposes to add oats to this cycle. But when 3 or 4 lines of oats are sown between the potatoes, it does not produce enough biomass to cover the soil during the whole year. The alternative of hairy vetch (*Vicia villosa*) needs water during the dry season, so it can only be used on irrigated plots. For the system to work, the sequence of crops needs to change: Maize together with potatoes and after that beans with oats. In this way, the oats grows enough to cover the soil and serve as a live cover crop. But at the moment, this sequence is not commonly practiced (Charpentier, 2010).

Table 0.1 Rotations at *tanety* in landratsay

	C1 : November - May	C2: April – October
Common system	Maize + Beans	Potato/Barley/Wheat
Proposed CA system	Maize + Potato	Beans + Oats/Vetch

VIEWS OF FARMERS

CROPS TO COVER OR TO USE ALTERNATIVELY

As noted before, the focus of this research is on the farm and village level. Here an important factor is the trade-off in the allocation of resources (CA2AFRICA, 2009). The fierce competition for crop residues, as described by Giller *et al.* (2009), also arises here. For a farmer, there are several possibilities when it comes to using his crop residues. He can use it as feed for his cattle, or as straw which mingles with the cow dung to become fertilizer. He can also burn it to obtain fertile ashes, or compost it. When a farmer implements a CA system, the possibility of leaving the residues on the field is added to these. Dairy production is an important component of the regional economy. And as land is scarce, every piece of vegetation is preferably used to feed the cows. Farmers even told that, when they decided to leave the plants on their field, someone else stole it (Ahmim-Richard *et al.*, 2010). This theft of crop residues is a real problem. It even occurs with living plants, which is the reason why farmers plant the cassava plants very close together. In this way, the roots become deep and difficult to pull out (oral information, O. Husson).

The high pressure on soil resources was also observed in Betafo by Kasprzyk (2008). The dairy farmers work with a “zero-loss” system, in which it is unimaginable to leave crop residues on the field. For these farmers, labour is not expensive so the labour-saving side of CA does not appeal to them. The systems that have potential should produce more biomass than these farmers are currently experiencing. The amount needed to cover the soil is about 6-7 ton/ha. BVPI SE/HP identifies this pressure on the use of crops as the main limiting factor for CA practices (Raharison & Andrianaivolala, 2009). In the CA systems, it is also possible to use living crops for coverage of the soil. Mulching of dead plants is better for the soil micro biota, but living cover crops have a higher potential for farmers (Rabary *et al.*, 2008). Quite surprising, the research of Razafimandimby (2007) found out that farmers prefer the dead mulch over living cover crops.

AREA AVAILABLE FOR AGRICULTURE

Especially in places with rich soils (like in landratsay), the population density is very high. This leads to small parcels, as shown in the figures in subsection 2.1.3. When a farmer has such a limited amount of land, it leads to relatively huge losses if he sets apart some land for cover crops. The *Brachiaria* needs for example a year of growth to be able to enrich the soil.

FINANCIAL SITUATION

In Antsapanimahazo , most of the farmers that have never tried CA, mentioned the high costs of investment in CA as an obstacle for implementing it. The CA systems were in fact sold as a package in the framework of intensification of agriculture. The systems require investment in chemical inputs and specific equipment. These investments will not pay back before several years, and in the first years the revenue from the parcel will be lower than before. For the farmers whose resources are very scarce, it is important to minimize risks. The access to credit does not resolve this, because the interest rates are said to be very high. When comparing different scales of income, it becomes clear that the capacity to bear risks is a determining factor for deciding about CA. Most of the farmers who have practiced CA for 3 years or more, are relatively rich (Randrianarison *et al.*, 2008). The lack of money was also the main problem that farmers mentioned in the research of Razafimandimby (2007), conducted in Antsapanimahazo, Ampandrotrarana and Ivory. TAFE offers no assurance if the harvest is lost, which can happen through natural causes. This research concluded that the credit system should become less rigid, to enable more farmers to profit from it. It also observed that rich landowners do not differ significantly from the average farmer in their willingness to practice CA.

Minimizing risk also often means that farmers prefer an off-farm job above their work at the farm. This does not automatically lead to abandoning CA practices, but Randrianarison *et al.* (2008) found that it did happen, especially when the father of the family was working off-farm. This leads to the question why their wives apparently decided to abandon CA. Households with a small amount of available labour and small capital availability to employ additional labourers, could profit from CA practices because it often requires less labour. But the figures on this are not very convincing. (Razafimandimby, 2007). Another important factor is the type of tenure. The poor farmers often rent the land, with contracts that are just valid for several years. This is another reason why long-term investments are probably not advantageous for them (Randrianarison *et al.*, 2008).

EXPERIENCE WITH THE PROJECT

Farmers that have never tried to implement CA, are often not enough informed about the system. Witnessing other people abandoning CA is also a reason to stay away from it.

In cases where people have tried CA but abandoned it after some time, the organisation of the dissemination turned out to be problematic. Credit can only be obtained when one is a member of a farmer's association. There is a lot of critique on these organisations. Complaints are about the delivery of inputs and material, that is often late (Randrianarison *et al.*, 2008).

CONCLUSION: HYPOTHESES

With this information from different resources, we return to the question: What are the most important factors that explain the hesitant attitude of farmers towards CA in this region? A summary of the constraints that follow from experiences of BVPI, SCRID, TAFA and research of farmers' preferences, in order of importance:

- The choice for cover crops is restricted due to the cold climate.
- Livestock feeding requirements compete with CA practices (mulch in the fields) for biomass.
- The small size of the farming enterprises leaves little room for risks or 'unproductive' cover crops.

Reasoning from these constraints, the hypotheses that are tested in the survey are the following:

1. The size of the farm (more land exploited, more cattle, more income) has a positive effect on adoption of CA.
2. When crops on paddy fields represent the big(gest) share of agricultural activity, this diminishes motivation for CA.
3. If the workload of women gets more heavy with CA practices, they are more inclined to decide against CA.
4. When deciding about adoption of CA, economic reasons are the determining factor.

Martin expects a change from CA

Martin Randrianandrasana (21 years old) and his wife (19) live in Ambohimarina, landratsay. They married in 2006 and have a son of 3 years old. They own 0.26 ha of land and also work off-farm on other people's land; the kind of off-farm employment that generates the lowest income (about 0.5 euro per day). They are resource-poor farmers. Last year they bought a pig, but had to sell it at a loss.

Since 2008, Martin has practiced CA on some land. He is not convinced of the benefits of the system; especially the maize did not grow and produce well. He finds it hard to keep up with the techniques of the system. Martin and his wife took part in the project with the expectation to enhance their standard of living. This has not yet become true.

PERSONAL STORIES II



Figure 5-1 Martin Randrianandrasana and his wife

Eliane has never practiced CA

Eliane Lalahine Rabemananjara (32) lives at Anjanamanjaka, Fitakimerina, with her husband (32) whom she married at the age of 17. She is a kind young woman and is surrounded by her 6 children during the interview. Except for the youngest two, all of them attend school. The production from their tiny agricultural enterprise (0.15 hectare, no cattle) does not match up with the food requirements of the family. This is why both parents work a lot for other farmers. It is clear that they belong to the type of 'resource-poor farmers'.

She is not familiar with the concept of CA but has practiced a kind of controlled underground burning. The system consists of digging a hole, burning residues in it and then install sweet potato. They are not really into contact with an extension agent, but saw someone else do it and found it productive.

Marie Henriëtte abandoned CA and the project

Rasoarimanana Marie Henriëtte (50) is married and has five children who all attend school. Their farm at Ampamelonana, landratsay, consists of 0.19 hectare of land and some poultry. 2 or 3 days a week, the parents work at other people's lands.

She is dissatisfied with the project organisation. According to her, the technician blocked the installation of her producers' organisation. Marie Henriëtte tried CA one time, in 2007. She has grown some wheat and vetch, but did not get seeds to continue. At the moment, she is not into contact with a technician.



Figure 5-2 Zebus at work while we interviewed Marie Henriëtte

Daniel has not yet tried CA

Daniel Rabenjamin (58) and his family migrated to Anjanamanjaka, Fitakimerina, in 2008. Daniel is retired from his job at the Ministry of Agriculture, where he supervised rural construction works. His wife (54) works as a seamstress. The pension provides for a substantial off-farm income. They have 7 children, of which 4 still live with them.

Their farm covers a relatively large area: 2.5 hectare. The work is mostly delegated, so they hire a lot of (wo)manpower. Daniel has heard of CA and is associated to the project BVPI SE/HP. He thinks that the characteristics of the soil are a determining factor for success of CA. But it would take a lot of training to actually practice the system. Since they have just arrived in the area, Daniel has not tried it yet, but he is interested to enrich his experience in agriculture, especially if this would increase production.

Results of survey

This paragraph will present the data obtained that describe the actual situation in the survey areas. These data cover the year from April 2009 – May 2010. Most of the data is grouped according to application of CA (practising, abandoned or never practiced), zone (Fitakimerina and landratsay, see subsection 2.1.1) and farm type (as described in section 4.3). Table 6.1 gives information about the composition of the distinguished groups. The distribution between zones and CA adoption is rather even; the same counts for the ratio of CA (non)adopters per farm type. The distribution of the farm types among the zones is less even. In both zones, about half of the surveyed farmers belongs to the farm type 'resource-poor'. One-third is receiving substantial off-farm income and a minority is of the 'producing' type. The presentation of the data will follow the outline of respectively: zone – farm type – CA adoption. Amounts of money are expressed in *Ariary*, the monetary unit of Madagascar with a rough exchange rate of 2600 *Ariary* per euro in July 2010.

Table 0.1 Number of farmers applying CA or not, by farm type and by zone

		Farm type			Zone		Total
		Off-farm	Producing	Poor	Fitakimerina	landratsay	
CA adoption	Practising	7	4	10	10	11	21
	Abandoned	5	3	9	9	8	17
	Never applied	6	4	12	11	11	22
Zone	Fitakimerina	10	5	15			30
	landratsay	8	6	16			30
Total		18	11	31			60

Households are most often composed of about 5 persons; this does not differ much between the groups. One of the strata was the gender of the interviewed farmer. During the survey, it turned out that in a married situation, the husband is practically always head of the household. Often during an interview with a woman, her husband joined in the conversation. It is also a cultural aspect, in which a woman does not decide about farming strategies, unless she is the single head of the family. This is why the statistics for age were taken for the male part of a married couple. Exceptions to this were two single women, of which one divorced and one widow. For labour availability, children that go to school are not considered. Those that are not at school are counted 0.5 if they are 10-15 years old. People older than 60 are also counted for 0.5 person equivalent.

Table 0.2 Family and labour features

Characteristic	Total	CA adoption			Zone		Farm type*		
		Pract	Aband	Never	Fitak	landr	Off	Prod	Poor
Household size (persons)	5	5	5	6	5	5	6	4	6
Age head of household (hoh)	44	39	44	48	45	43	45	41	44
Education level hoh**	1.5	1.6	1.3	1.6	1.6	1.5	1.6	1.7	1.4
Labour availability within family (person equivalent)	2.1	1.8	2.2	2.4	2.2	2.1	1.8	2.3	2.3
Off-farm employment of household (days)	158	153	146	171	125	191	323	33	107
Off-farm income of household (1000 Ariary)	871	889	859	863	822	919	2 180	205	347
Number of hired man days	83	92	64	90	63	103	85	126	67

*) For explanation of these abbreviations, see section 4.3

***) Value between 1 = read and write, 2 = finished secondary school, 3 = finished university

Table 6.2 shows that people in landratsay count more days of off-farm employment. Numbers of hired man days are also higher here. The other factors are more equally distributed. Labour availability within family shows no clear pattern. The off-farm employment and income are highest with the farm type 'Off-farm', as can be expected. Off-farm income ranges largely between zero and 1 800 thousand, with 6 exceptional cases up to 6 000 thousand Ariary. Concerning the age of the head of the household, one thing is interesting: for the group that is practising some form of Conservation Agriculture, average age is lowest. This while their education level is the same as that of those who never practiced CA. For the group that has abandoned CA practices, the number of hired man days is very low.

The survey also paid attention to tasks that are usually done by women. This was done in a qualitative way. Most common activities that are assigned to women are sowing of different crops, replanting of the rice and transport of manure and fertilizer.

EQUIPMENT

Angady is a Malagasy term for a digging stick with a blade at its end. It is a tool that is very generally present. The spraying equipment was also part of the survey and turned out to be scarcely present. Only in Fitakimerina some people (9) own it; usually it is a shared tool from the organisation of FIFAMANOR. This originally Norwegian organisation has in the past introduced dairy cattle and is still mainly active in the field of milk and forage, but also with the dissemination of other crops.

Table 0.3 Equipment features

Equipment	Total	CA adoption			Zone		Farm type		
		Pract	Aband	Never	Fitak	landr	Off	Prod	Poor
Percentage of households that owns this equipment									
<i>Angady</i> (kind of spade)	97	100	94	95	97	97	89	100	100
Weeder	65	76	53	64	73	57	56	91	61
Plough	37	38	41	32	40	33	6	27	23
Cart	18	19	24	14	13	23	28	64	32
Average number of tools per household									
<i>Angady</i>	2.8	2.5	3.0	2.8	2.5	3.0	2.4	3.3	2.8
Weeder	0.9	1.0	0.8	1.0	1.1	0.7	0.8	1.6	0.7
Plough	0.2	0.2	0.2	0.1	0.1	0.2	0.1	0.3	0.2
Cart	0.4	0.4	0.4	0.3	0.4	0.3	0.3	0.6	0.3
Average number of tools for households that own this equipment									
<i>Angady</i>	2.9	2.5	3.2	3.0	2.6	3.1	2.7	3.3	2.8
Weeder	1.4	1.3	1.4	1.5	1.5	1.3	1.4	1.8	1.2
Plough	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
Cart	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0

Only people who are receiving substantial off-farm income do not always own an *angady*. For the other equipment it is clear in table 6.3 that the full-time farmers ('producing' type) have the highest figures. Distribution between CA (non) adopters is very homogeneous.

LIVESTOCK

The figures in table 6.4 only present the adult animals. The animal unit that is used counts Zebus and cows as 1, pigs as 0.2 and poultry as 0.008 per unit.

Table 0.4 Livestock features

Type of livestock	Total	CA adoption			Zone		Farm type		
		Pract	Aband	Never	Fitak	landr	Off	Prod	Poor
Animal unit	1.75	2.08	1.54	1.54	1.37	2.12	1.78	1.98	1.54
Liters of milk sold	314	166	421	372	67	561	408	300	245
Percentage of households that own this kind of livestock									
Zebu	35	52	29	23	37	33	33	64	26
Cow (female)	42	43	35	45	30	53	39	27	48
Pig	57	57	35	73	53	60	50	73	55
Poultry	87	86	94	82	80	93	78	82	94
Average number of animals per household									
Zebu	0.7	1.0	0.6	0.5	0.7	0.7	0.7	1.5	0.5
Cow (female)	0.7	0.8	0.6	0.7	0.4	1.0	0.8	0.5	0.7
Pig	1.0	0.9	0.9	1.1	0.7	1.3	1.0	1.0	1.0
Poultry	14.6	17.5	15.2	11.2	8.2	20.9	17.1	7.6	15.6
Average number of animals for households that own this type of livestock									
Zebu	2.0	1.9	2.2	2.2	1.9	2.2	2.0	2.3	1.9
Cow (female)	1.7	1.8	1.7	1.6	1.4	1.8	2.0	1.7	1.5
Pig	1.7	1.6	2.6	1.5	1.3	2.1	2.0	1.3	1.8
Poultry	16.8	20.5	16.1	13.7	10.3	22.4	22.0	9.2	16.6

It is visible that in landratsay the emphasis is a bit more on dairy cattle, although the higher figure for milk sales also comes from the overall difference in prosperity of the zones. This will also be dealt with in the next paragraphs. Within the farm types, the 'off-farm' type sells the highest amount of milk. The group of people that practises CA has the highest average animal unit. This can be contributed to the relatively high percentage of households that own Zebus in this group.

LAND AND CROPS

Table 6.5 presents the data obtained about owned land. According to the interviewed farmers, there is no formal tenure system. People usually inherit their land, and buy some more. Only in landratsay, 9 people hired land during the surveyed year. Sharecropping and lending were less observed. For paddy fields, the number of women involved in planting was used to calculate the area. This results in most cases in a totally different figure than the estimation of the interviewed farmer, but it was considered more accurate. The estimation of BVPI is that 1 woman can plant 1.5 are. Paddy fields turned out to be practically always in use. Only one farmer mentioned 20 are of fallow paddy fields, because of inundation. A remark should be made here about the term for 'paddy fields' that was used during the survey – i.e. the Malagasy word *tanimbary* (*tany*= land, *vary*= rice). It could be that this word is interpreted by the farmers in the strict sense of a rice field. In this case, inundated fields are not considered to be *tanimbary*. For the total area of land used per person, every member of the family counts as 1, except babies. Measuring the importance of paddy fields relative to *tanety*, turned out to be complicated. When asked, almost everyone in Fitakimerina mentioned the paddy fields as their first priority. In landratsay, the answers were more diffuse. Although rice is the main staple crop, products from *tanety* fields provide cash income and some poor farmers only have land at *tanety*. The input invested in both types of fields was surveyed by a very general question – if or not fertilizer (NPK and/or urea) had been used. Table 6.5 shows the average scores.

Table 0.5 Land features

Feature	Total	CA adoption			Zone		Farm type		
		Pract	Aband	Never	Fitak	landr	Off	Prod	Poor
Age of farm (years until 2010)	18	15	19	20	19	18	19	15	18
Soil type*	2.35	2.44	2.20	2.39	1.90	2.81	2.46	1.95	2.37
Area paddy fields (are)	28	31	22	29	34	22	30	52	16
Area <i>tanety</i> fields (are)	94	57	159	76	53	133	66	190	68
Percentage of <i>tanety</i> fields that are fallow (are)	27	30	28	22	34	19	23	38	23
Total area of used land per person (are)	38	16	98	13	12	64	14	143	10
Percentage that applied fertilizers to paddy fields	63	76	65	50	57	70	56	58	68
Percentage that applied fertilizers to <i>tanety</i>	65	90	76	32	53	77	72	50	65

*) Code: 1=sandy, 2=ferrous, 3=volcanic

Apparently, people in landratsay do not only dispose of the largest area of *tanety* fields but also actively use the largest percentage of it. Fertilizers are more generally used than in Fitakimerina. The soil is also more fertile in landratsay. Overall, the full-time farmers ('producing' type) have a relatively large farm area. On average, the CA-practising farmers run the 'youngest enterprises'. They also slightly more often apply fertilizers to their fields. Of the people that have never applied CA, only a third has used fertilizers for *tanety* cropping. Figures that stand out, are the high average area of *tanety* fields used and of area per person, of the group that has abandoned CA practices.

Table 0.6 Most occurring and CA cropping patterns on *tanety*, number of farmers practising (first column) and average area used (in are)

Cropping pattern*	No.	Average area (are)	CA adoption			Zone		Farm type		
			Pract	Aband	Never	Fitak	landr	Off	Prod	Poor
(Potato/) Maize + [beans or soja]	39	24	26	33	15	7	33	14	28	27
CA Maize+[beans or soja] /oats	4	5	5	5	-	-	5	5	3	5
CA Potato+oats / maize+beans	1	3	3	-	-	-	3	-	-	3
CA Vetch / maize+beans	1	15	15	-	-	-	15	-	-	15
Sweet potato(+maize) / upland rice	3	38	10	4	100	7	100	4	10	100
CA Sweet potato+maize / upland rice+stylo	1	2	2	-	-	2	-	-	2	-
Brachiaria	9	9	9	6	11	12	8	4	7	13
CA Cassava+brach / cassava(+maize)+beans	2	7	7	-	-	7	-	-	13	2
CA Brachiaria / beans+soja	1	3	3	-	-	3	-	-	-	3
CA Brach/sw.pot/upl.rice +soja	1	2	2	-	-	2	-	-	-	2
Cassava	11	7	13	7	5	5	8	1	8	6
Beans or soja	12	7	-	10	4	4	9	5	6	12

*) Between () is optional, between [] is a choice. After / comes the next season.

CA cropping systems are mostly practiced on the *tanety* fields, and so were the described CA cases of this survey. That is why table 6.6 covers only those fields. The figures for average area are only

calculated from farmers who practice the particular cropping pattern. A hectare of sweet potato followed by upland rice from a farmer in landratsay is very visible in the table. The table describes 11 cases of CA that were surveyed in detail. From the remaining 10 CA practising farmers, no detailed information was available. One pattern is clear – the CA systems that involve oats are only practiced in landratsay, while those with Brachiaria were only found in Fitakimerina. Because of the low numbers, it is not possible to draw conclusions from this.

Farm income

Table 0.7 Average production sold (kg/year)

Crop	Total	CA adoption			Zone		Farm type		
		Pract	Aband	Never	Fitak	landr	Off	Prod	Poor
Potato	306	257	429	257	29	583	491	309	187
Barley	257	324	165	265	83	431	89	377	301
Rice	90	96	61	108	124	57	38	267	50
Maize	39	21	69	32	4	73	31	58	35
Beans	34	52	26	25	3	66	49	28	28
Soja	32	33	51	16	1	64	52	4	30
Wheat	8	9	0	14	0	16	14	17	1
Sweet potato	8	12	6	5	15	0	0	37	1

Table 6.7 shows the quantities of crops that are traded most frequently. Except for one rich, female farmer in Fitakimerina who sold 2500 kg, barley is only cultivated in landratsay (see subsection 2.1.3). The amounts in table 6.8 represent just the gross income from sales of agricultural products, including animals (mostly poultry and pigs) and milk. It can serve as an indicator of the relative importance of trade between the groups. It is clear that production in landratsay is much more market-oriented than it is in Fitakimerina. Between CA (non) adopters there is not really a clear pattern.

Table 0.8 Returns from agricultural products (1000 Ariary)

	Total	Application of CA			Zone		Farm type		
		Pract	Aband	Never	Fitak	landr	Off	Prod	Poor
Gross income from sales of agricultural products	718	754	686	707	174	1 261	785	748	643
Gross income (as above) divided by total area in use	15	19	9	14	7	22	18	16	12

Overall picture

The data described above firstly show the difference between the two zones. In landratsay there exists more of a money economy, a fact that is already clear when one notices children with the newest models of cell phones. People also work more off-farm than in Fitakimerina. The number of animals per household is higher in landratsay, except for Zebus. This zone also has the largest area of tanety fields per person. The CA systems that are practiced are rotations with oats, while in Fitakimerina this is Brachiaria or Stylosanthes (1 time).

Secondly, the description of the different farm types generally meets the characteristics that were laid out in section 4.3. The ‘producing’ type hires the most labour and has the highest average area of tany fields per person. In paddy fields, the difference is not significant. The ‘off-farm’ type generally owns less equipment, but sells the highest amount of milk. Regarding crops, they mainly sell potatoes.

Table 0.9 Correlation between farm features and CA adoption (P=2, A=1, N=0)

	Spearman’s rho	Significance 2-tailed	Significance 1-tailed
Age	-0.341**	0.008	
No. Zebus	0.233*	0.073	0.037
Input paddy	0.230*	0.077	0.038
Age of farm	-0.186	0.155	0.078
Income sold crops	0.119	0.364	0.182
Area pp	0.073	0.578	0.289
Off-farm income	-0.043	0.742	0.371

*) significant at 0.05 level

***) significant at 0.01 level

Lastly and most interesting for this research, there are some factors that distinguish between the different groups of CA adoption. Factors of resources seem of little importance in this regard. Table 6.9 presents the results of a statistical analysis of correlation between farm features and CA adoption. The only outstanding figures are the high number of Zebus and a higher use of fertilizers on paddy fields for the group that practices CA, which are not very significant. Farmers who abandoned CA practices have a large average tany area but the smallest paddy area, although this last difference is not significant. The only factor that correlates firmly to adoption of CA is a lower age of the head of household. This does not seem to have a link with education level.

SOCIAL FEATURES INFLUENCING ADOPTION

The BVPI SE/HP project and its operator SDMAD work through producers’ associations and technicians in the fields. Table 6.10 lists some of the features that describe people’s experience with the project.

Table 0.10 Experience with CA project

Feature	Total	CA adoption			Zone		Farm type		
		Pract	Aband	Never	Fitak	landr	Off	Prod	Poor
% that knows the system (explains it right)	62	86	82	23	67	57	72	73	52
Knowledge of CA test field*	0.9	1.8	0.5	0.5	0.4	1.5	1.1	0.9	0.8
% that is connected to producers’ association	58	95	88	0	63	53	67	55	55
No. of years CA practiced	2	4	2	0	2	2	2	2	2
Known number of people who practice CA	4.5	6.3	0.9	5.8	4.1	4.9	4.3	9.3	2.7
Known number of people who have abandoned CA	9.1	10.3	14.3	4.0	7.7	10.5	10.2	9.2	8.2

*) Code between 0 = not heard of, 1 = knows but not visited, 2 = visited with project, 3 = visited ‘voluntary’

Not everyone that practices CA describes this system as a no-tillage rotation with cover crops. Within the definition of the FAO, the emphasis lies on covering the soil. During the survey, no one mentioned rotation as part of CA, and they often viewed mulching in combination with tillage as CA. As can be expected, with the farmers who never practiced CA, knowledge of the system is lowest. The test fields are not widely known. The producers' associations have a strong link with the extension of CA practices. Not one of the farmers that never applied CA is connected to such an association. It is clear that CA is still in a start-up phase. Farmers have only very recently started trying CA practices (since 2006). No one practices it on a significant scale (see table 6.6).

In general, people know more people who have abandoned CA than people who practice it. The important influence of social structures becomes clear in the example of Tsaratanana (Fitakimerina), where the whole family of the technician is involved in the project. Other people feel ignored, and this does not stimulate a good relationship between the technician and potential CA practisers.

DECIDING ABOUT APPLICATION OF CA

One of the pillars of Conservation Agriculture is the absence of tillage. Since this is a concept that is very unfamiliar to the surveyed farmers, it is interesting to understand their view on tillage. Table 6.11 lists the reasons for tillage.

Table 0.11 Percentage of farmers that mentioned these reasons for tillage (more than one reason per farmer)

Reason for tillage	Total	CA adoption			Zone		Farm type		
		Pract	Aband	Never	Fitak	landr	Off	Prod	Poor
Necessary for production	70	67	59	82	60	80	61	73	74
Soil structure	20	14	29	18	27	13	33	27	10
Out of habit	12	14	18	5	7	17	6	18	13
Soil fertility	8	14	6	5	3	13	6	18	6
To control weeds	7	5	6	9	3	10	11	0	6

When asked about their primary reason to plough the soil, most people answered that it is necessary for production. Surprisingly, this number is lowest with farmers who abandoned CA practices. The question that followed was why it is necessary. These reasons are all interrelated, but table 6.11 attempts to categorize them somewhat. Maintaining/restoring a soil structure that favours plant growth is mentioned most often, especially in Fitakimerina. Maintaining soil fertility (by decomposition of OM and a flush of mineralized nitrogen) and controlling weed growth are in general functions of tillage (Wall, 2007). In this case, most farmers do not perceive it as a primary reason to plough. Table 6.12 presents the different reasons that were mentioned to practice or not practice CA. For analysis, they were grouped in four categories: economical, soil features, professional development in agriculture and influences from social environment. In general, people in landratsay mention slightly more positive arguments than in Fitakimerina. After the table, the categories are described in more detail.

Table 0.12 Percentage of farmers that mentioned this reason to practice (+) or not practice (-) CA (more than one reason per farmer)

Category of reason		Total	CA adoption			Zone		Farm type		
			Pract	Aband	Never	Fitak	landr	Off	Prod	Poor
Economical	+	37	52	18	36	23	50	44	33	32
	-	48	48	71	32	50	47	61	33	45
Soil features	+	20	29	6	23	20	20	17	50	10
	-	20	14	29	18	27	13	22	8	23
Professional development in agriculture	+	12	10	6	18	13	10	22	0	10
	-	15	14	18	14	13	17	11	8	19
Influences social environment	+	13	14	0	23	7	20	11	17	13
	-	28	10	41	36	30	27	33	25	26

The **economic** (dis)advantages of the system are mentioned most often. Figures 6-1 and 6-2 show the distribution of these arguments in more detail, with figures that can also be found in Appendix C.




Figure 0-1 Number of farmers that mentioned economic advantages of CA

A higher production and smaller demand for labour were the main economic advantages (figure 6-1). These figures, as well as the percentage for lower expenses, are mainly shaped by opinions of CA practising farmers. The factor ‘subsidised agricultural inputs’ was mostly mentioned by the group that has never applied CA practices.




Figure 0-2 Number of farmers that mentioned disadvantages of CA

It is in particular the opinion of the group that has abandoned CA practices, that contributes (with 9 farmers) to a ‘low production’ (figure 6-2). This is also the main economic disadvantage according to both this group and the practising group. ‘Time and labour consuming’ was mentioned evenly by the

three groups. With regard to soil features, the advantage of CA is the improvement of the soil quality. Negative arguments are that the system is not adapted to the local soil which according to the farmers is either very hard to cultivate on or too poor. Reasons to practice CA that were mentioned in the sphere of professional development in agriculture were to gain experience, test the system and get instructed in general. The fact that the system is too difficult to follow is seen as a disadvantage. The influences from the social environment can be positive: the examples from others and the encouragement of extension staff. On the other hand there are also negative examples of people who abandoned CA that form a reason against the system. This negative influence of the social environment is mentioned more often and includes social or personal constraints to enter a producers' group and complaints about extension staff. People that have never practiced CA, have significantly little to say about the production level of the system. In general, their positive and negative arguments weigh up against each other, except for influences from the social environment that tend towards a negative result.

Table 0.13 Experienced and expected impact of CA systems for three parameters: score from -1 (lower) to 1 (higher)

Parameter	Practising: experienced	Abandoned: Experienced	Never applied: expected
Availability of fodder	0.75	0.13	0.90
Harvest	0.14	-0.65	0.17
Expenses	-0.67	0.06	-0.60

Table 6.13 presents the view of farmers on the impact of practising (a) CA system(s). The availability of fodder is both experienced and expected to increase; the cover crops involved in CA rotations are really perceived as a provision of fodder. Expenses become smaller with CA, mainly because of the less hired labour that is needed to plough the soil. According to these figures, farmers that abandoned CA do not find lower expenses with CA, and are also dissatisfied about the production of the system. Interestingly, the people that never applied CA have high expectations for both the availability of fodder, the increase of their harvest and the decrease of their expenses. Regarding the experiences, at least the first two expectations seem to be too optimistic.

DISCUSSION

Reasoning from the data presented above, this paragraph will discuss the hypotheses that were formulated at the end of chapter 5. The size of the farm (more land exploited, more cattle, more income) has a positive effect on adoption of CA. With regard to the area of the farm, this hypothesis can be rejected. Farmers who abandoned CA have the largest area per person. The relation could be the other way around, if a large cropping area discourages the adoption of time-consuming rotations. However, the reasons in table 6.11 do not explicitly point towards this.

The amount of cattle does not differ significantly between the groups of CA application, with the exception of Zebu. About half of the practising farmers owns (a) Zebu(s), while with the other groups this is less than 30%. The fact that the number of Zebus does at least not correlate to a lower application of CA, seems to indicate that the competition of cover crops and cattle does not occur within a farm. This can be explained by an observation from the project, that residues are also sold (Raharison & Andrianaivolala, 2009).

The income from crops sales is slightly higher for people who practice CA, but this is not significant. Comparison with off-farm income also shows no correlation; contrary to conclusions of Clay *et al.* (1998), availability of money does not stimulate adoption of CA. So overall, this hypothesis was only confirmed with regard to the amount of Zebus. Another observation with a stronger relation, was that lower age of the head of household correlates positively to adoption of CA. This outcome is the opposite of earlier mentioned research in Ethiopia (de Graaff *et al.*, 2008). Age can both stimulate and hinder changes in agricultural practices; in this case younger farmers are apparently more motivated to use CA than their older colleagues. Education level has no relation with this.

When crops on paddy fields represent the big(gest) share of agricultural activity, this diminishes motivation for CA.

Comparing the importance of *tanety* and paddy fields turned out to be difficult, especially with a general survey that does not include detailed information about labour inputs. The measured factor was input spent on the fields in terms of fertilizers. At the paddy fields, this does not differ a lot between the CA (non) adopters. However, there is a pattern whereby practising farmers apply most and those who never tried CA least. This leads to a tentative rejection of the hypothesis.

In general, paddy will always stay more important relative to *tanety*. Rice is a source of liquidity; saved amounts are sold when people need money to repair their house or to pay salaries.

If the workload of women gets more heavy with CA practices, they are more inclined to decide against CA.

When a woman is married, it is her husband who makes the decisions. During most of the interviews that were arranged with these women, her husband also took part. Estimations of workloads for women would be biased by this. This hypothesis can be rejected because in general women do not decide about implementation of CA.

When deciding about adoption of CA, economic reasons are the determining factor.

The effect of CA on production levels and demand for labour is definitely an important factor. Complaints about a bad production and positive experiences with economized labour expenses are the two opposite sides of the coin. Those people that have no experience with CA, apparently do not decide on the basis of production levels but see benefits in the subsidised inputs that the project provides. The perception of many of these farmers is that production will not increase with a CA system. An often heard remark was that “the soil here is too hard and tillage is necessary to have a production.” These are all economic reasons. But apart from this, it also seems that CA practices are extension-intensive. Farmers think it will not work without a lot of attention from SDMA-employees. Social constraints form the second largest group of negative arguments. For a farmer to jump the train of this new agricultural system, it is vital to be part of a producers’ association. This implies good contacts with co-farmers and extension staff from BVPI SE/HP, and that is exactly what is lacking in many cases.

Lastly, this discussion treats the information from the 11 farmers who described a CA system that they practiced (introduced in table 6.6 and described in more detail in Appendix C). The five systems that involve oats as a cover crop are all located in landratsay. This could be related to the relatively higher importance of dairy production at this location. These systems still combine maize and beans

like in the usual rotation. Three of these farmers want to continue CA on that location. A weakness seems to be the sensitivity to uncontrolled growth of weeds. The farmers that do not want to continue, find the production of the system (with regard to maize and beans) lower than it usually would be. If production would not be influenced by the combination with oats, people would probably adopt it more. The four farmers that implemented a CA system with *Brachiaria* were all content with the production levels and wanted to continue. The problem of weeds was not mentioned. It is clear that they practice it on a very small scale.

Discussion of methods

FIGURES FOR THE FARM TYPOLOGY

To use the farm typology of Ahmim-Richard *et al.* (2010), a figure for 'self sufficiency in rice' had to be calculated. Because of an initial different understanding of this term, the total production of rice was not asked in the survey. In order to apply to the definition of BVPI, this figure had to be calculated from the area of paddy fields and the average yield, which does not make it very precise.

A situation of self sufficiency in rice is defined as: The household eats rice for at least 8 months a year. It is calculated as follows:

$$\frac{\text{Number of are paddy} \times \text{Average yield per are}}{\text{Number of persons at charge} \times \text{Per capita rice consumption for eight months}}$$

In this way, an outcome ≥ 1 indicates that the household is self sufficient in rice.

The per-capita consumption of rice is estimated at 250 kg paddy/year. Because the survey did not include the difference between well- and poorly managed parcels (RMME, *rizières à mauvaise maîtrise de l'eau*), the average yield of the paddy fields is a very rough estimate. Rakotofiringa *et al.* (2007) mention an average yield for irrigated rice of 3.3 ton/ha for the village of Andranomanelatra (north of Antsirabe). It appears that this counts for the well-managed paddy fields. In the research of Ahmim-Richard *et al.* (2010), figures vary according to the different types of farmers: 2 (type 1), 2.5 (type 5), 3 (type 4) or 3.2 (type 3) ton/ha. BVPI SE/HP (2009a) mentions 3 ton/ha for irrigated rice (well-managed) in the highlands but has no figures for RMME in this region. In the South-East region it ranges from 0.53 to 1.72 ton/ha. It is not clear which part of the paddy fields can be qualified as RMME. The location of Fitakimerina reports 30% of the paddy-fields well-irrigated and 70% poorly irrigated (BVPI SE/HP, 2008a), but this does not necessarily mean that 70% of the fields are RMME. In general, the area of landratsay seems to have no RMME at all (BVPI SE/HP, 2009a).

Considering all this, the average yield of the paddy fields was estimated to be 2.5 ton/ha in landratsay and 2 ton/ha in Fitakimerina. To account for losses during harvest and storage (10%) the figures that are used are 22.5 kg/are and 18 kg/are. FAO data for Madagascar show a comparable yield of 2.5 t/ha in 2008 (FAOSTAT, 2010).

A last remark on the farm typology: It is said (Ahmim-Richard *et al.*, 2010) that type 5 occurs only in the middle-west part of Vakinankaratra and Amoron'i Mania, but seeing that in this survey 6 cases fell into this category, type 5 is also present in the highlands.

Social dimension

In the process of determining the sample, the technicians were too influential. As indicated in chapter 4, they were asked to find people to interview within the strata of gender, farm size and number of dairy cows. In this way, the survey in Fitakimerina was mainly held in the sublocation of Anjanamanjaka. Only the last day was spent in Tsaratanana, a location closer to the highway and because of that with some different features. A better distributed sample would have resulted in a more balanced image. Everyone who has never practiced CA, said he or she wanted to start with it. This is most probably caused by their perception of the interview – we were seen as extension agents who can provide them with seeds and inputs. It is difficult, even impossible, to judge whether people tell their true thoughts about CA or whether they tell a story to receive the things that they want to get. In general, it is preferred to use a research method that avoids a one-way flow of information, from the researcher to the subject. It is a bit suspicious that the research team requires knowledge of a person's age, income and possessions, while they themselves do not reveal information about these personal details. At least the purpose of the survey should be clear, ideally in a way that makes the results (this report) available to the interviewed farmers.

SOURCES OF INACCURACY

The survey dangled a bit between an in-depth study to farming systems and a qualitative research to opinions about CA. It would have been better to start with an empty database and create a survey to collect the data. There are also constraints in data collection that probably every farming system survey encounters. Most of the time, the area of parcels is not readily known. The survey was not designed to visit all the fields, so this led to estimates like '2 times that field over there'. The same counts for questions about soil types. If it is not possible for the researcher to visit the fields, it is better to take pictures (or samples) of defined soil types and make it a multiple choice.

3 CONCLUSIONS

"On ne développe pas, on se développe"
Prince Claus of the Netherlands (1926-2002)

This research aimed at identifying the past and present reasons why farmers are hesitant to adopt Conservation Agriculture practices in the highlands of Vakinankaratra. In the first place, one should note that real adoption, in the sense of continued use of CA practices, is not yet possible because the introduction of CA in these zones only started in 2006. According to the results of the survey, the only factor that correlates firmly with adoption of CA is a lower age of the head of household. The amount of Zebus has a slightly positive effect on adoption of CA, so competition for livestock fodder seems no big hindrance. Clearly, it is also not a matter of resources - relatively within these zones, that is to say. Of course the surveyed farmers all have small enterprises and live in an agricultural society with little resources. The idea of CA does not raise high expectations with regard to production. This is mainly due to a lack of confidence in a no-tillage system; *'labour toujours'* seems to be the device in the studied zones. People who applied CA mention this low production level as the main economic disadvantage. Farmers without experience with CA are more inclined to decide on the basis of subsidised inputs they can get through the project. The social threshold they need to take to get involved in the project is their weakest link towards CA. This shows that the local context involves more than 'just' climatic circumstances and financial possibilities. Changing an agricultural

practice requires strong support systems that provide inputs and equipment (Corbeels *et al.*, 2011), and above all a social environment that incites and stimulates this change.

Elaborating on the notion of change - the meaning of this concept depends highly on a person's circumstances. While a Dutch student may find moving to another country a big change, for a farmer in Fitakimerina a change in crop rotation has much more influence on the income of the family and issues like food security. So, one of the reasons why farmers are 'hesitant' to apply CA practices, could be that they simply have no choice. Several of the families we encountered during the survey had no capital to invest in whatever better system. While they are the ones that can use innovation of agricultural practices, they are caught in the poverty-trap and do not have any power to choose. 'Development' in this sense means that they themselves create a way to raise their production or income. For future study, it would be interesting to compare the dissemination of CA to another (past) project of SDMAD in the region: reforestation with Eucalyptus. During the survey it became clear that it was more popular than CA. Many trees could be seen and farmers mentioned it as a way to protect the soil and the environment. Apart from questions about the effects of Eucalyptus, there could be a lesson with regard to adoption.

Other examples are for instance the organisation of Malto, who buys barley from farmers (see subsection 2.1.3) and the dairy producing industry that stimulates dairy cattle. In these cases, there exists a market for the products. When de Graaff *et al.* (2008) conclude that profitability is an important stimulus for investment in conservation practices, a special niche for CA-products can be given some thought.

References

AFD/FFEM (2007) *Direct seeding mulch-based cropping systems (DMC)*. Paris, France.

Ahmim-Richard, A., Bodoy, A., Penot, E. (2010). *Caractérisation et typologie des exploitations agricoles dans le Vakinankaratra et l'Amoron'i Mania, Madagascar*. Document de travail BV lac n° 25BVPI/SCRID/FOFIFA/TAFA.

BVPI SE/HP (2008a) *Diagnostic terroir Ambohimanga-Tsaratana (Commune Vinaninkarena, Region Vakinankaratra, Lot 1)*. Version du 16-06-08.

BVPI SE/HP (2008b) *Diagnostic terroir Ampamelomana (Périmètre landratsay, Commune Mandritsara, Region Vakinankaratra)*. Version du mars 2008.

BVPI SE/HP (2008c) *Diagnostic terroir Est Anosy (Périmètre landratsay, Commune Mandritsara, Region Vakinankaratra)*. Version du mars 2008.

BVPI SE/HP (2009a) *Rapport Tri Annuel : octobre 2006 – septembre 2009*.

BVPI SE/HP (2009b) *Rapport de campagnes, region Vakinankaratra, Saison 2008-2009*.

BVPI SE/HP (2010a) *Base des données Vakinankaratra*, Microsoft Access document.

BVPI SE/HP (2010b) *Evolution de surface en SCV durant les 4 années du projet*.

CA2AFRICA (2009) *Annex I – Description of work*. Conservation Agriculture in AFRICA: Analysing and Foreseeing its Impact - Comprehending its Adoption.

Critchley, W. (ed.) (1999) *Promoting Farmer Innovation Workshop Report no 2*. UNDP/SIDA.

Charpentier, H. (2010) *Rapport de mission auprès du projet BVPI SE/HP, 28 avril au 8 mai 2010*.

CIRAD (2010) *Website ‘Le Cirad à Madagascar’*, URL: www.cirad.mg [29-03-2010].

Clay, D., Reardon, T., Kangasniemi, J. (1998) Sustainable Intensification in the Highland Tropics: Rwandan Farmers’ Investments in Land Conservation and Soil Fertility. *Economic Development and Cultural Change*, 46 (2): 351-377.

Corbeels M., Apina T., Koala S., Schuler J., Triomphe B., El Mourid M., Traore K., Nyagumbo I., Mrabet R., Penot E., Gomez-Macpherson H., de Graaff J., Tiftonell P. (2011) *Impact and adoption of conservation agriculture in Africa: a multi-scale and multi-stakeholder analysis*.

FAO (2009) Adoption of Conservation Agriculture Technologies: Constraints and Opportunities, in: *Proceedings of the 4th World Congress on Conservation Agriculture, Lead Papers*, New Delhi, 4-7 February, 257-264.

FAO (2010) *Website ‘Conservation Agriculture’*. URL: www.fao.org/ag/ca [21-04-2010].

FAOSTAT (2010) *Website ‘FAOSTAT’*. URL: <http://faostat.fao.org> [2-11-2010].

Fowler, R., Rockström, J. (2001) Conservation tillage for sustainable agriculture—an agrarian revolution gathers momentum in Africa. *Soil & Tillage Research*, 61, 93–107.

Giller, K.E., Witter, E., Corbeels, M., Tiftonell, P. (2009) Conservation agriculture and smallholder farming in Africa: The heretics’ view. *Field Crops Research*, 114, 23-34.

Gowing, J.W., Palmer, M. (2008) Sustainable agricultural development in sub-Saharan Africa: the case for a paradigm shift in land husbandry. *Soil Use and Management*, 24, 92–99.

Graaff, J. de, Amsalu, A., Bodnár, F., Kessler, A., Posthumus, H., Tenge, A. (2008) Factors influencing adoption and continued use of long-term soil and water conservation measures in five developing countries. *Applied Geography*, 28, 271-280.

Hobbs, P.R. (2007) Conservation agriculture: what is it and why is it important for future sustainable food production? *Journal of Agricultural Science*, 145, 127–137.

Hobbs, P.R., Sayre, K., Gupta, R. (2008) The role of conservation agriculture in sustainable agriculture. *Philosophical Transactions of the Royal Society B*, 363, 543–555.

Kasprzyk, M. (2008) *Diversité des systèmes d’alimentation des troupeaux bovins laitiers à Betafo, Région du Vakinankaratra, Madagascar*. Thesis.

Knowler, D., Bradshaw, B. (2007) Farmers’ adoption of conservation agriculture : A review and synthesis of recent research. *Food Policy*, 32, 25-48.

Rabary, B., Sall, S., Letourmy, P., Husson, O., Ralambofetra, E., Moussa, N., Chotte, J. (2008) Effects of living mulches or residue amendments on soil microbial properties in direct seeded cropping systems of Madagascar. *Applied Soil Ecology*, 39, 236-243.

Raharison, T., Andrianaivolala, E. (2009) *Rapport Tri Annuel Octobre 2006 à Septembre 2009, Volet Agro Ecologie*, Cellule de projet BVPI SE/HP.

Rakotofiringa, A., Tokarski, Y., Penot, E. (2007) *Caractérisation des exploitations agricoles dans la commune rurale d'Andranomanelatra, Région Vakinankaratra, hauts plateaux de Madagascar*. Thesis.

Randrianarison, Narilala, Penot, Eric, Poncet, Christian (2008), *Suivi et analyse des succès et des abandons des systèmes à base de semis direct sous couverture végétale (SCV) : mise au point de la méthodologie. Cas du fokontany d'Antsapanimahazo – Madagascar*. Document de travail BV lac n° 3

Razafimandimby, A.J.W, Razafimandimby, S. (2007) *Etude des mécanismes d'adoption des SCV : cas d'Antsapanimahazo, Ampandrotrarana et d'Ivory, Dans la région du Vakinankaratra*. Document de travail BV lac n° 7, BVPI/SCRID/FOFIFA/TAFA.

TAFA-FIFAMOR (1995) *Rapport de campagne 1994-1995*.

Seguy, L. (2003) *Les techniques de semis direct sur couvertures végétales à Madagascar, ou comment pratiquer une agriculture durable avec un minimum d'intrants chimiques. Le cas des régions des Hauts Plateaux*.

Wall, P.C. (2007) Tailoring Conservation Agriculture to the Needs of Small Farmers in Developing Countries. *Journal of Crop Improvement*, 19 (1), 137-155.

APPENDIX A – DETERMINATION KEYS FOR FARM TYPOLOGY




Figure A-1 Determination keys for farm typology (Source : Ahmim-Richard *et al.*, 2010)

